





Remove the cap from the mixing liquid.

2


- Keep the vial stable.
- Push one of the spikes through the • membrane of the vial.




- Attach the purple syringe to the spike. Withdraw saline only up to the red mark on the
- syringe, there will be saline left in the glass vial.
- Disconnect the saline-filled syringe from the spike.




Push the second spike through the membrane of the gentamicin vial. Attach the saline-filled syringe to the spike.




- Dissolve all the gentamicin powder.




Retract the blue handle and remove the red plunger stop ring.







- b) Attach valve, clear end to the powder-filled syringe, and turn clockwise.
- c) Attach syringe with gentamicin solution to blue end of valve, press and turn clockwise.




- Detach the syringe from the valve.



- After the liquid has been injected, start the stopwatch immediately.
- Mix with rotation for 30 seconds, at 1 complete stroke per second.
- Push the handle all the way down and pull it all the way up, and rotate at its end positions.



Fully retract blue handle. a) b) Lock plunger by turning blue collar clockwise to click.



CERAN 🔨

CERAN

 \mathbf{igodol}

- Attach dispensing syringe to valve. a)
- b) Immediately transfer paste with numbers facing user. When full, paste will begin to ooze from under sleeve. Stop filling when this occurs.
- to use.

back into the syringe. This may require the spike to be slightly backed out.

Inject all the gentamicin solution into the syringe with powder. Keep the plunger pushed to the bottom to avoid back flush.



Remove the red stopper and the paste is now ready

ADDITIONAL RESOURCES: Review our mixing video here:





PR 01326-01 en AU CA EU ROW 04-2023